

Posttraumatic stress disorders in children and adolescents

Bruce D. Perry, MD, PhD*, and Ishnella Azad†

Millions of children are exposed to traumatic experiences each year. Over 30% of these children develop a clinical syndrome with emotional, behavioral, cognitive, social, and physical symptoms called posttraumatic stress disorder. The symptoms of posttraumatic stress disorder fall into three clusters: reenactment of the traumatic event; avoidance of cues associated with the event or general withdrawal; and physiological hyperreactivity. Significant physical and medical problems in childhood, adolescence, and adulthood appear to be related to childhood trauma. Current treatment approaches include postacute psychoeducation, individual psychotherapy, pharmacotherapy, and cognitive-behavioral therapy. Despite increasing attention over the past 10 years, childhood posttraumatic stress disorder remains an understudied public health problem. *Curr Opin Pediatr* 1999, 11:310–316 © 1999 Lippincott Williams & Wilkins, Inc.

*CIVITAS Academy & Departments of Psychiatry, Pediatrics, Pharmacology and Neuroscience, Baylor College of Medicine, Houston, Texas, USA. †MS IV, Baylor College of Medicine, Houston, Texas, USA

Correspondence to Bruce D. Perry MD, PhD, Texas Children's Hospital, The Feigin Center, Suite 715, 6621 Fannin Street, Houston, TX 77030-2399, USA; e-mail: bperry@bcm.tmc.edu

Current Opinion in Pediatrics 1999, 11:310–316

Abbreviations

DSM IV *Diagnostic and Statistical Manual of Mental Disorders*, fourth edition
PTSD posttraumatic stress disorder
PTSS posttraumatic signs or symptoms

ISSN 1040-8703 © 1999 Lippincott Williams & Wilkins, Inc.

Neurophysiological core of posttraumatic stress disorder

Each year in the United States more than five million children are exposed to some form of extreme traumatic stressor. These traumatic events include natural disasters (*eg*, tornadoes, floods, hurricanes), motor vehicle accidents, life-threatening illnesses and associated painful medical procedures (*eg*, severe burns, cancer), physical abuse, sexual assault, witnessing domestic or community violence, kidnapping, and sudden death of a parent, among others [1,2••]. These events, posing an actual or perceived threat to the individual, activate a stress response. During the traumatic event, the child's brain orchestrates adaptive stress-mediating neural systems including the hypothalamic-pituitary-adrenal axis, central nervous system noradrenergic, and dopaminergic systems and associated central and peripheral nervous system mechanisms that provide the adaptive emotional, behavioral, cognitive, and physiological changes necessary for survival [3•].

Individual adaptive responses during traumatic stress are heterogeneous [4,5]. The specific nature of a child's responses to a given traumatic event may vary with the nature, duration, and pattern of traumatic stressor and the child's constitutional characteristics (*eg*, genetic predisposition, age, gender, history of stress exposure, presence of attenuating factors such as supportive caregivers). Whatever the individual response, however, the extreme nature of the external threat is often matched by an extreme and persisting internal activation of the neurophysiological systems mediating the stress response and their associated functions [3•]. A primary adaptive feature of the threat-response system is single-trial "learning"—the capacity to generalize from a threatening event to other experiences with similar features. Unfortunately, this very adaptive capacity is at the core of the emotional, behavioral, and physiological symptoms that develop following a traumatic experience.

Neural systems respond to prolonged, repetitive activation by altering their neurochemical and, sometimes, microarchitectural (*eg*, synaptic sculpting) organization and functioning. This is no different for the neural systems mediating the stress response. Following any traumatic event, children will likely experience some persisting emotional, behavioral, cognitive, and physiological signs and symptoms related to the sometimes

temporary shifts in their internal physiological homeostasis. In general, the longer the activation of the stress-response systems (*ie*, the more intense and prolonged the traumatic event), the more likely there will be a “use-dependent” change in these neural systems [3•,4,6]. In some cases, then, the stress-response systems do not return to the pre-event homeostasis. In these cases, the signs and symptoms become so severe, persisting, and disruptive that they reach the level of a clinical disorder [5]. In a new context and in the absence of any true external threat, the abnormal persistence of a once-adaptive response becomes maladaptive.

Posttraumatic stress-related clinical syndromes

Posttraumatic stress disorder (PTSD) is a clinical syndrome that may develop following extreme traumatic stress *Diagnostic and Statistical Manual of Mental Disorders [DSM IV]* [7]. Like all other *DSM IV* diagnoses, it is likely that heterogeneous pathophysiologies underlie the cluster of diagnostic signs and symptoms labeled PTSD. With this in mind, there are six diagnostic criteria for PTSD: extreme traumatic stress accompanied by intense fear, horror, or disorganized behavior; persistent reexperiencing of the traumatic event such as repetitive play or recurring intrusive thoughts; avoidance of cues associated with the trauma or emotional numbing; persistent physiological hyperreactivity or arousal; signs and symptoms present for more than 1 month following the traumatic event; and clinically significant disturbance in functioning. A child is considered to have acute stress disorder (*DSM IV*) when these criteria are met during the month following a traumatic event. PTSD is further characterized as acute when present for less than 3 months, chronic when present for more than 3 months, or delayed onset when symptoms develop initially 6 months or more after the trauma.

Posttraumatic stress disorder has been studied primarily in adult populations, most commonly, combat veterans and victims of sexual assault. Despite high numbers of traumatized children, the clinical phenomenology, treatment, and neurophysiological correlates of childhood PTSD remain understudied. The clinical phenomenology of trauma-related neuropsychiatric sequelae are poorly characterized [8,9]. Most of the studies of PTSD have been following single discrete traumas (*eg*, a shooting). The least-characterized populations are very young children and children who have undergone multiple or chronic traumatic events.

Several factors complicate the study of PTSD in children. It has only been in the past 10 years that child-specific, structured interviews for PTSD have become available. The development of trauma-specific psycho-

metrics continues [10,11]. In very young children, diagnostic assessment is difficult due to the inability of infants and toddlers to self-report trauma-related symptoms, the differential expression of symptoms across the developmental spectrum, and the difficulty determining the nature and extent of certain traumatic experiences (*eg*, exposure to domestic violence or physical abuse) [12,13]. A key complication in studying and treating trauma-related neuropsychiatric problems in children is the complex and varied clinical presentations that may result following acute or chronic trauma [8].

Clinical presentation

Children with PTSD may present with a combination of problems including impulsivity, distractibility, and attention problems (due to hypervigilance), dysphoria, emotional numbing, social avoidance, dissociation, sleep problems, aggressive (often reenactment) play, school failure, and regressed or delayed development. In most studies examining the development of PTSD following a given traumatic experience, twice as many children suffer from significant posttraumatic signs or symptoms (PTSSs) as are diagnosed with PTSD; however, children diagnosed as having PTSSs lack all of the criteria necessary for the diagnosis of PTSD [14]. In these cases, the clinician may identify the trauma-related symptom as being part of another neuropsychiatric syndrome.

The clinician is often unaware of ongoing traumatic stressors (*eg*, domestic or community violence), or the family makes no association between the present symptoms and past events (*eg*, car accident, death of a relative, exposure to violence) and may provide no relevant history that would aid the clinician in the differential diagnosis. As a result, PTSD is frequently misdiagnosed, and PTSSs are underrecognized. Children with PTSD as a primary diagnosis are often labeled with attention deficit disorder with hyperactivity, major depression, oppositional defiant disorder, conduct disorder, separation anxiety, or a specific phobia. Ackerman *et al.* [15•] examined the prevalence of PTSD and other neuropsychiatric disorders in 204 abused children (ages 7 to 13 years). Thirty-four percent of these children met criteria for PTSD. More than 50% of the children in this study who had suffered both physical and sexual abuse had PTSD. Using a structured diagnostic interview, the majority of these children met diagnostic criteria for three or more *DSM IV* Axis I diagnoses in addition to PTSD. Indeed, only six of 204 children met criteria for only PTSD. The broad comorbidity reported in this study echoes previous studies.

Diagnostic criteria from the *DSM IV* yield multiple labels in maltreated children, but these diagnoses rarely provide useful information about etiology, course, treat-

ment response, or prognosis. At present, despite an evolving clinical phenomenology, it is clear that PTSD is not the only, nor an inevitable, outcome of traumatic events during childhood. Posttraumatic hyperarousal or dissociative-like symptoms often coexist with these other axis I disorders. Furthermore, severe early trauma appears to be an expresser of underlying constitutional or genetic vulnerability and may be a primary etiologic factor in the development of a broad range of disorders later in life.

Incidence and prevalence

Estimates of lifetime incidence of PTSD range from 3% to 14% [7]. Cuffe *et al.* [16•] examined population prevalence of PTSD in a community sample of adolescents. They found that 3% of girls and 1% of boys met the *DSM IV* criteria for PTSD. In this study, girls reported more traumatic events than boys. Being female, experiencing rape or sexual abuse, and witnessing an accident or medical emergency were associated with increased risk for PTSD. Children exposed to various traumatic events have much higher incidence (from > 15% to > 90%) and prevalence rates than the general population [1]. Several studies published in 1998 confirm previous reports of high prevalence rates for PTSD in high-risk groups. Thirty-five percent of a sample of adolescents in whom cancer was diagnosed met criteria for lifetime PTSD [17•]; 15% of children surviving cancer had moderate to severe PTSS [18]; 93% of a sample of children witnessing domestic violence had PTSD [19]; over 80% of the Kuwaiti children exposed to the violence of the Gulf Crisis had PTSS [20]; 73% of juvenile male rape victims develop PTSD [21]; 34% of a sample of children experiencing sexual or physical abuse and 58% of children experiencing both physical and sexual abuse all met criteria for PTSD [15•]. In all of these studies, clinically significant symptoms, though not full PTSD, were observed in essentially all of the children or adolescents following the traumatic experiences.

Vulnerability and resilience

Not all children exposed to traumatic events develop PTSD. A major research focus has been identifying factors (mediating factors) that are associated with increased risk (vulnerability) or decreased risk (resilience) for developing PTSD following exposure to traumatic stress [19]. Factors previously demonstrated to be related to risk can be summarized in these broad categories: characteristics of the child (*eg*, subjective perception of threat to life or limb, history of traumatic exposures, coping style, general level of anxiety, gender, age); characteristics of the event (*eg*, nature of the event, direct physical harm, proximity to threat, pattern and duration); characteristics of the

child's family or social system (*eg*, supportive, calm, nurturing versus chaotic, distant, absent, anxious) [18,22,23]. Each of these mediating factors can be related to the degree to which they either prolong or attenuate the child's stress-response activation resulting from the traumatic experience. Factors that increase stress-related reactivity (*eg*, family chaos) will make children more vulnerable, whereas factors that provide structure, predictability, nurturing, and sense of safety will decrease vulnerability. Persistently activated stress-response neurophysiology in the dependent, fearful child will predispose to "use-dependent" changes in the neural systems mediating the stress response, thereby resulting in posttraumatic stress symptoms.

Adolescents with cancer who developed PTSD rated their families as more chaotic than adolescents with cancer who did not develop PTSD [17•]. Most interesting in this study, however, was that 85% of mothers of the PTSD group also developed PTSD related to their child's cancer. If the family is chaotic and the primary caregiver is traumatized by an event, their capacity to provide a consistent, predictable, and nurturing environment is compromised.

There are apparent gender differences in the expression and development of PTSD. Clinical experience and recent studies suggest that girls tend to exhibit more internalizing (*ie*, anxiety, dysphoria, dissociation, avoidance) and boys more externalizing (*ie*, impulsivity, aggression, inattention, hyperactivity) posttraumatic symptoms [4,15•]. In epidemiologic studies of PTSD in the general adult population, women have higher rates than men [24]. Although the extensive epidemiologic data from these adult studies are lacking in studies of children, a gender difference has been observed in several studies with children and adolescents [15•]. There appear to be gender differences in adaptive response in the acute event (girls dissociate more than boys) that may be related to this observed difference in development and expression of trauma-related symptoms [4].

Long-term consequences of childhood trauma

Posttraumatic stress disorder is a chronic disorder. Untreated, PTSS and PTSD remit at a very low rate. Indeed, the residual emotional, behavioral, cognitive, and social sequelae of childhood trauma persist and appear to contribute to a host of neuropsychiatric problems throughout life [25••], including attachment problems [26,27], eating disorders [28], depression [23,25••], suicidal behavior [29], anxiety [25••], alcoholism [25••,30], violent behavior [25••,31], mood disorders [32] and, of course, PTSD [33,34].

Childhood trauma affects other aspects of physical health throughout life as well [35•,36]. Adults victimized by sexual abuse in childhood are more likely to have difficulty in childbirth [37], a variety of gastrointestinal and gynecologic disorders, and other somatic problems such as chronic pain, headaches, and fatigue [37]. The Adverse Childhood Experiences study [38••] examined exposure to seven categories of adverse events during childhood (*eg*, sexual abuse, physical abuse, witnessing domestic violence: events associated with increased risk for PTSD). This study found a graded relationship between the number of adverse events in childhood and the adult health and disease outcomes examined (*eg*, heart disease, cancer, chronic lung disease, and various risk behaviors). With four or more adverse childhood events, the risk for various medical conditions increased four- to 12-fold.

Special concerns for pediatrics

Pediatricians should be aware that children with PTSD or PTSS may have altered sensitivity and functioning of neuroendocrine and autonomic nervous systems [6,35•,39–41]. This altered sensitivity may predispose to the development of various medical conditions such as asthma, hypertension, cardiac arrhythmias, endocrine disorders, gastrointestinal disorders, and various other somatic complaints [36]. Furthermore, PTSD complicates the treatment of various medical conditions [42]. In children with diabetes, for example, the PTSD-related hyperreactivity of the counterregulatory hormones such as adrenaline may complicate or prevent effective control of blood sugar. History of sexual or physical abuse can complicate the medical examination of traumatized children, manifesting as resistance to medical examination and procedures [43,44]. In a variation of PTSS adaptation, excessive compliance, “numbing,” and insensitivity to pain may also be seen in children with histories of chronic exposure to traumatic violence in the home. Addressing posttraumatic stress symptoms within a multidisciplinary approach is an important component of improved outcomes following childhood injuries [45].

Treatment approach

To date, few treatment outcome studies in children with PTSS and PTSD have been published. Despite this dearth of objective data, a wealth of papers on clinical experience and subjective treatment approaches have been published [1]. The nature of these reported clinical approaches depends on the theoretical perspective of the author. At present, the mechanism-based conceptual frameworks explaining the development of PTSD fall into four main categories: 1) psychoanalytic, 2) cognitive-behavioral, 3) psychodevelopmental, and 4) neurodevelopmental. Each of these offers certain insights but none provides a complete and unambiguous

treatment approach. Therefore, the treatment of children with PTSD varies greatly depending on the specific clinician’s training, perspective, and experience. Typically, the nature and severity of specific symptoms (*eg*, impulsivity, withdrawal, hypervigilance, dissociation, dysphoria, and aggression) dictate treatment approach rather than the diagnosis. Another major consideration in treatment is distinguishing between a single discrete traumatic event (*eg*, car accident or witnessing an assault) and chronic or pervasive trauma (*eg*, chronic abuse). Symptoms following a single event (*eg*, motor vehicle accident) tend to be fewer and less treatment resistant compared with the more complex symptom clusters associated with chronic or pervasive traumatic stress (*eg*, a combination of physical and sexual abuse). A host of clinical treatments are used with traumatized children, including family therapy, group therapy, eye-movement desensitization and reprogramming (EMDR), music and movement therapies, “play” therapy, and art therapy, among many others. Four of the major therapeutic approaches used alone or in combination are discussed here.

Secondary prevention using acute posttraumatic interventions

In the immediate posttraumatic period, several models of intervention have been used to diminish the acute distress and improve posttraumatic outcome [46]. One of the most important is psychoeducation. Telling the family and child what the expected signs and symptoms are following a traumatic event can help diminish anxiety, increase sense of competence, and provide a baseline from which parents and children can be aware of abnormally intense or prolonged symptoms requiring further clinical attention. Several modifications of a critical incident stress-debriefing paradigm have been reported, though efficacy has not yet been determined. In some cases, clinicians have used antianxiety agents or clonidine to decrease the level of physiological hyperarousal and distress in the acute posttraumatic period [47]. Although these postacute interventions are clinically helpful during this period, it is not yet clear that any of them alter the development, course, or severity of PTSD.

Pharmacotherapy

There have been very few published trials of psychotropic medications in childhood PTSD [1,5]. Without the benefit of clinical outcome studies, the selection of psychotropic agents has been guided by empirical clinical judgment and the clinical observations that primary symptoms in PTSD appear to respond to psychotropic agents proven to be useful for those symptoms in other neuropsychiatric disorders (*eg*, valproic acid and lithium for aggressive behavior, fluoxetine for depressive symptoms).

Many of the symptoms of PTSD can be traced to the core symptoms of physiological hyperarousal such as sleep problems (including difficulties falling asleep, early night awakening, nightmares, night terrors), generalized anxiety, behavioral impulsivity, or hyperreactivity of the sympathetic nervous system including tachycardia, hypertension, increased muscle tone, respiratory problems, and body temperature dysregulation. Clonidine, an α_2 -adrenergic partial agonist that modulates the reactivity of the locus caeruleus and decreases the physiological hyperreactivity associated with PTSD, has been shown to be an effective agent in children with PTSD [6]. Other agents altering the biogenic amines (*ie*, serotonin, dopamine, and norepinephrine) may also modulate the symptoms of PTSD. In this regard, preliminary reports support the efficacy of propranolol and fluoxetine in children with anxiety and PTSD [5].

Individual psychotherapy

The core hyperarousal symptoms result in a cascade of secondary, interrelated problems. Inability to engage in appropriate intimacy leads to difficulties with peer and adult relationships, and inability to perform adequately in school leads to poor self-esteem, resulting in a variety of learned behaviors that both mask and defend against these core deficits driven by their physiological hyperreactivity. The resulting vicious cycle of poor performance, poor self-esteem, and development of maladaptive problem-solving styles, in turn, is difficult to treat as long as the underlying physiological hyperreactivity impairs the ability to modulate anxiety, concentrate on academic or social learning tasks, and contain behavioral impulsivity. Successful treatment, therefore, often requires "containing" or modifying this core physiological dysregulation with medications and using other psychotherapeutic interventions to address issues related to self-esteem, competence, social skills, and mastery of specific fears.

Cognitive-behavioral therapies

Cognitive-behavioral therapy is the most studied and likely the most effective therapeutic intervention in adults with single-event-related PTSD. The few studies of cognitive-behavioral therapy in children and adolescents are very promising [48]. March *et al.* [49••] examined a standard cognitive-behavioral therapy protocol in school-aged children following a single traumatic event. After the course of treatment, significant improvement was noted in all main dependent measures. Unfortunately, cognitive-behavioral therapy is difficult to apply in the same fashion to very young children or to children with chronic pervasive trauma.

Conclusions and future directions

Despite the progress of the past few years, childhood PTSD remains a woefully understudied disorder.

Conservative estimates of the frequency of traumatic events (> five million children traumatized per year) and the well-documented incidence rates of more than 30% following a trauma suggest that there may be as many as 1.5 million children developing PTSD each year. Further, based on the documented incidence from high-risk populations, another 1.5 million may have clinically significant posttraumatic stress symptoms that do not meet full PTSD criteria. PTSD and PTSS are chronic problems. Available data show only moderate rates of remitted symptoms over time; in contrast, adolescents and adults who have undergone childhood trauma appear to be more vulnerable to a host of medical and neuropsychiatric problems. Better characterized clinical phenomenology, outcome studies examining a variety of therapeutic modalities, and mechanism-focused neurophysiological studies are necessary to better characterize PTSD and the other sequelae of childhood trauma.

Acknowledgments

The authors would like to acknowledge the support of CIVITAS Initiative, The Hogg Foundation for Mental Health, Children's Crisis Care Center of Harris County, the Ella T. Fondren Trust, and the Azzam Foundation.

References and recommended reading

Papers of particular interest, published within the annual period of review, have been highlighted as:

- Of special interest
- Of outstanding interest

- 1 Pfefferbaum B: Posttraumatic stress disorder in children: a review of the past 10 years. *J Am Acad Child Adolesc Psychiatry* 1997, 36:1503-1511.
- 2 Pfefferbaum B: Stress in Children. In *Child Adolesc Psychiatr Clin N Am*, •• vol 7. Edited by Lewis M. Philadelphia: W.B. Saunders; 1998. This contributed volume summarizes the current state of clinical, research, and policy-related issues in the area of childhood traumatic stress. Several of the primary theoretical constructs guiding research and treatment are outlined. Excellent summaries of clinical experience and reviews of current clinical research are also included.
- 3 Perry BD, Pollard R: Homeostasis, stress, trauma, and adaptation: a neurodevelopmental view of childhood trauma. *Child Adolesc Psychiatr Clin N Am* 1998, 7:33-51. This review examines the available neurodevelopmental and neurophysiological studies related to childhood trauma. The authors revise previously stated neurodevelopmental theoretical constructs that had been used to guide clinical research and practice. This synthesis focuses on memory and the neural systems involved in the stress response.
- 4 Perry BD, Pollard RA, Blakley TL, Baker WL, Vigilante D: Childhood trauma, the neurobiology of adaptation and use-dependent development of the brain: how states become traits. *Infant Ment Health J* 1995, 16:271-291.
- 5 Perry BD: Anxiety disorders. In *Textbook of Pediatric Neuropsychiatry*. Edited by Coffee CE, Brumback RA. Washington: American Psychiatric Press, Inc.; 1998:580-594.
- 6 Perry BD: Neurobiological sequelae of childhood trauma: post-traumatic stress disorders in children. In *Catecholamines in Post-traumatic Stress Disorder: Emerging Concepts*. Edited by Murberg M. Washington: American Psychiatric Association Press, 1994:253-276.
- 7 American Psychiatric Association: *Diagnostic and Statistical Manual of Mental Disorders*, edn 4 (DSM IV). Washington: American Psychiatric Association Press; 1994.
- 8 Terr L: Childhood traumas: an outline and overview. *Am J Psychiatry* 1991, 148:1-20.
- 9 Mulder RT, Fergusson DM, Beautrais AL, Joyce PR: Relationship between dissociation, childhood sexual abuse, childhood physical abuse, and mental illness in a general population sample. *Am J Psychiatry* 1998, 155:806-811.

10 Kent A, Waller G: The impact of childhood emotional abuse: an extension of the Child Abuse and Trauma Scale. *Child Abuse Negl* 1998, 22:393-399.

11 Matorin A, Lynn SJ: The development of a measure of correlates of child sexual abuse: the traumatic sexualization survey. *J Trauma Stress* 1998, 11:261-280.

12 Scheeringa MS, Zeanah CH, Drell MJ, Larrieu JA: Two approaches to the diagnosis of post-traumatic stress disorder in infancy and early childhood. *J Am Acad Child Adolesc Psychiatry* 1995, 34:191-200.

13 Scheeringa MS, Zeanah CH: Symptom expression and trauma variables in children under 48 months of age. *Infant Ment Health J* 1995, 16:259-270.

14 Friedrich WN: Behavioral manifestations of child sexual abuse. *Child Abuse Negl* 1998, 22:523-531.

15 Ackerman PT, Newton JE, McPherson WB, Jones JG, Dykman RA: Prevalence of post traumatic stress disorder and other psychiatric diagnoses in three groups of abused children (sexual, physical, and both). *Child Abuse Negl* 1998, 22:759-774.

This study examined PTSD and other neuropsychiatric disorders in over 200 maltreated children. This study used excellent, structured interviewing methods for diagnostic assessment. Although the total sample was small, this study is important because of the rigor used in determining comorbid diagnoses. Of interest is the demonstration of the symptoms and outcome differences between physical and sexual abuse, the increased risk with both types of abuse, and the gender differences in trauma-related outcomes.

16 Cuffe SP, Addy CL, Garrison CZ, Waller JL, Jackson KL, McKeown RE, Chilappagari S: Prevalence of PTSD in a community sample of older adolescents. *J Am Acad Child Adolesc Psychiatry* 1998, 37:147-154.

This study is the second cycle of a longitudinal epidemiologic study. In this cycle the authors examined a population sample of 490 adolescents (age range, 16 to 22 years) and used a semistructured interview to elicit PTSD symptoms and related factors. Of interest was the demonstration of a gender difference in the prevalence of PTSD (girls 3%, versus boys 1%). Being raped, witnessing a medical emergency, and witnessing an accident were associated with increased risk for developing PTSD. In this study, most of the children experiencing a traumatic event developed PTSD.

17 Pelcovitz D, Libov BG, Mandel F, Kaplan S, Weinblatt M, Septimus A: Posttraumatic stress disorder and family functioning in adolescent cancer. *J Trauma Stress* 1998, 11:205-221.

This study compared 23 adolescents with cancer against 27 physically abused and 23 healthy, nonabused adolescents. Of primary interest was the rate of lifetime PTSD (35% of patients in the cancer group, compared with only 7% of patients in the abused group). In the PTSD-positive subgroup of children, 85% of the mothers also developed PTSD. This study is very important for practicing pediatricians. The rate of PTSD in life-threatening pediatric illness is high for both the child and for caregivers. This has profound implications for creating a multidimensional clinical approach to the treatment of children with cancer.

18 Stuber ML, Kazak AE, Meeske K, Barakat L, Guthrie D, Garnier H, et al.: Predictors of posttraumatic stress symptoms in childhood cancer survivors. *Pediatrics* 1997, 100:958-964.

19 Kilpatrick KL, Williams LM: Potential mediators of post-traumatic stress disorder in child witnesses to domestic violence. *Child Abuse Negl* 1998, 22:319-330.

20 Hadi FA, Llabre MM: The Gulf crisis experience of Kuwaiti children: psychological and cognitive factors. *J Trauma Stress* 1998, 11:45-56.

21 Ruchkin VV, Eisemann M, Hagglof B: Juvenile male rape victims: is the level of post-traumatic stress related to personality and parenting? *Child Abuse Negl* 1998, 22:889-899.

22 Briggs L, Joyce PR: What determines post-traumatic stress disorder symptomatology for survivors of childhood sexual abuse? *Child Abuse Negl* 1997, 21:575-582.

23 Winje D, Ulvik A: Long-term outcome of trauma in children: the psychological consequences of a bus accident. *J Child Psychol Psychiatry* 1998, 39:635-642.

24 Breslau N, Davis GC, Andreski P, Peterson EL, Schultz LR: Sex differences in posttraumatic stress disorder. *Arch Gen Psychiatry* 1997, 54:1044-1048.

25 Fergusson DM, Horwood LJ: Exposure to interparental violence in childhood and psychological adjustment in young adulthood. *Child Abuse Negl* 1998, 22:339-357.

This is a report from an 18-year longitudinal study of a birth cohort of 1,265 New Zealand children. Retrospective reports of exposure to interparental violence were obtained as well as a host of measures of mental, social, physical, antisocial, and criminal behavior. The adolescents and adults reporting the highest levels of exposure were at the greatest risk for mental health problems,

substance abuse, and criminal offending. This study is well conceived and the methods are very sound. The value of this study is in demonstrating the multiple adverse sequelae of domestic violence. The pervasive nature of domestic violence, and the recurring issues of "how damaging" exposure to interparental violence is, will be addressed by studies of this sort. Exposure to domestic violence may be as potentially traumatic and abusive as physical or sexual abuse.

26 Bell D, Belicki K: A community-based study of well-being in adults reporting childhood abuse. *Child Abuse Negl* 1998, 22:681-685.

27 Alexander PC, Anderson CL, Brand B, Schaeffer CM, Grelling BZ, Kretz L: Adult attachment and long-term effects in survivors of incest. *Child Abuse Negl* 1998, 22:45-61.

28 Rorty M, Yager J: Histories of childhood trauma and complex post-traumatic sequelae in women with eating disorders. *Psychiatr Clin North Am* 1996, 19:773-791.

29 Allen JR, Heston J, Durbin C, Pruitt DB: Stressors and development: a reciprocal relationship. *Child Adolesc Psychiatr Clin N Am* 1998, 7:1-18.

30 Epstein JN, Saunders BE, Kilpatrick DG, Resnick HS: PTSD as a mediator between childhood rape and alcohol use in adult women. *Child Abuse Negl* 1998, 22:223-234.

31 O'Keefe M: Posttraumatic stress disorder among incarcerated battered women: a comparison of battered women who killed their abusers and those incarcerated for other offenses. *J Trauma Stress* 1998, 11:71-85.

32 Molnar BE, Shade SB, Kral AH, Booth RE, Watters JK: Suicidal behavior and sexual/physical abuse among street youth. *Child Abuse Negl* 1998, 22:213-222.

33 Schaaf KK, McCanne TR: Relationship of childhood sexual, physical and combined sexual and physical abuse to adult victimization and posttraumatic stress disorder. *Child Abuse Negl* 1998, 22:1119-1133.

34 Ford JD, Kidd P: Early childhood trauma and disorders of extreme stress and predictors of treatment outcome with chronic posttraumatic stress disorder. *J Trauma Stress* 1998, 11:743-761.

35 Orr SP, Lasko NB, Metzger LJ, Berry NJ, Ahern CE, Pitman RK: Psychophysiological assessment of women with posttraumatic stress disorder resulting from childhood sexual abuse. *J Consult Clin Psychol* 1998, 66:906-913.

This investigative team has pioneered study of trauma-related neurophysiological changes using standard psychophysiological methods. In this study, 29 women with chronic PTSD following childhood sexual abuse showed greater physiological responses (heart rate, skin conductance, electromyographic findings) than women experiencing sexual abuse but not having PTSD. This responsivity was specific to the conditions involving sexual imagery and was not seen in the stressful, nonabuse-related situation. These preliminary studies illustrate some of the physiological hyperactivity that may underlie some of the documented, long-term medical and physical problems following childhood trauma. Studies such as these are required to elaborate mechanism-related models of trauma-related neuropsychiatric and medical problems.

36 Hertzman C, Wiens M: Child development and long-term outcomes: a population health perspective and summary of successful interventions. *Soc Sci Med* 1996, 43:1083-1095.

37 Rhodes N, Hutchinson S: Labor experiences of childhood sexual abuse survivors. *Birth* 1994, 21:213-220.

38 Felitti VJ, Anda RF, Nordenberg D, Williamson DF, Spitz AM, Edwards V, et al.: Relationship of childhood abuse and household dysfunction to many of the leading causes of death in adults: the adverse childhood experiences (ACE) study. *Am J Prev Med* 1998, 14:245-258.

This study was conducted by mailing questionnaires about adverse childhood experiences to 13,494 adults in a large health maintenance organization. The response rate was 70.5%. The responses were studied along with the results of a standard medical evaluation and measures of adult risk behavior, health status, and related issues. At least half of the respondents reported at least one, and more than one fourth of the respondents reported more than two, categories of adverse childhood experiences. A graded relationship between the number of categories of childhood exposure and the high-risk behaviors and diseases was demonstrated. This study reinforces the observations of many other studies using different methods and drawing on different specific childhood stressors. The relationships between "health" throughout the life cycle and stress or distress during development are very strong.

39 De Bellis MD, Chrousos GP, Dorn LD, Burke L, Helmers K, Kling MA, et al.: Hypothalamic-pituitary-adrenal axis dysregulation in sexually abused girls. *J Clin Endocrinol Metab* 1994, 78:249-255.

40 De Bellis MD, Lefter L, Trickett PK, Putnam FW: Urinary catecholamine excretion in sexually abused girls. *J Am Acad Child Adolesc Psychiatry* 1994, 33:320-327.

- 41 Stein MB, Yehuda R, Koverola C, Hanna C: Enhanced dexamethasone suppression of plasma cortisol in adult women traumatized by childhood sexual abuse. *Soc Biol Psychiatry* 1997, 42:680-686.
- 42 Chadwick DL: Medical consequences of child sexual abuse: commentary. *Child Abuse Negl* 1998, 22:551-552.
- 43 Berkowitz CD: Medical consequences of child sexual abuse. *Child Abuse Negl* 1998, 22:541-550.
- 44 Britton H: Emotional impact of the medical examination for child sexual abuse. *Child Abuse Negl* 1998, 22:573-579.
- 45 Hanfling M, Perry BD, Kozinetz C, Gill A, Tilbor A, Brams M, Levin H: Improved medical and psychosocial outcomes of injured children with multidisciplinary versus conventional follow-up. *Proceedings of the Fourth World Conference on Injury Prevention and Control 1998*. Amsterdam: Fourth World Conference on Injury Prevention and Control; 1998:28.
- 46 Pynoos RS, Goenjian AK, Steinberg AM: A public mental health approach to the postdisaster treatment of children and adolescents. *Child Adolesc Psychiatr Clin N Am* 1998, 7:195-210.
- 47 Famularo R, Kinscherff R, Fenton T: Propranolol treatment for childhood post-traumatic stress disorder, acute type. *Am J Dis Child* 1988, 142:1244-1247.
- 48 Deblinger E, Mcleer S, Henry D: Cognitive behavioral treatment for sexually abused children suffering post-traumatic stress. *J Am Acad Child Adolesc Psychiatry* 1990, 5:747-752.
- 49 March JS, Amaya-Jackson L, Murray MC, Schulte A: Cognitive-behavioral psychotherapy for children and adolescents with posttraumatic stress disorder after a single-incident stressor. *J Am Acad Child Adolesc Psychiatry* 1998, 37:585-593.
- This study tested a group-administered cognitive-behavioral treatment protocol with a single case across time and setting design. The children ($n = 17$) were selected from two elementary and two junior high schools and screened for single-event-related PTSD. Neuropsychiatric symptoms were measured using state-of-the-art instruments. Fourteen of the 17 children completed treatment. Significant improvement was observed, such that 57% no longer met diagnostic criteria for PTSD. Despite the small numbers, this is one of the few well-designed and controlled treatment outcome studies in the area of childhood PTSD.